

HIGHLIGHTS AND BREAKTHROUGHS

Making a fine-scale ruler for oxide inclusions

DONGZHOU ZHANG^{1,*}

¹Hawai'i Institute of Geophysics and Planetology, University of Hawai'i at Manoa, Honolulu, Hawaii 96822, U.S.A.

Constraining the phase relations and valence state of iron-bearing oxides is crucial to understanding the chemistry of Earth's mantle. In this issue of *American Mineralogist*, Uenver-Thiele et al. (2017a) present an experimental study on the phase diagrams of magnesioferrite-magnetite solid solutions at high pressures and high temperatures. By analyzing the compositions of the quenched products, Uenver-Thiele et al. were able to constrain the phase diagram of the $(\text{Mg}, \text{Fe}^{2+})\text{Fe}_3^{3+}\text{O}_4$ series, and they identified several new phases with non-conventional stoichiometry. From the phase diagrams of $(\text{Mg}, \text{Fe}^{2+})\text{Fe}_3^{3+}\text{O}_4$ spinels determined in this study and the stability fields of the new phases, the authors proposed an empirical method to recover the petrological history of magnesium-iron oxide inclusions in natural diamonds.

Oxygen and iron are the two most abundant elements in the Earth, and their compounds, the iron oxides, are common in the crust and the mantle. Iron is the most important multivalent transitional metal in the mantle, and the stability fields of different iron oxides covers the mantle conditions in the P - T - f_{O_2} space. Certain iron oxides, such as magnetite (Fe_3O_4) and wüstite (Fe_{1-x}O) and their solid solutions with periclase (MgO), have been used as scales to constrain the P - T - f_{O_2} history of petrological assemblages. One application of such an "oxide scale" is to determine the petrological history of diamonds with inclusions. For example, diamonds with ferropericlase $[(\text{Mg}, \text{Fe})\text{O}]$ inclusions are usually associated with a lower mantle origin (Wirth et al. 2014), whereas magnesioferrite (MgFe_2O_4) inclusions have a maximum formation pressure of 8–10 GPa (Uenver-Thiele et al. 2017b). The oxygen fugacity of petrological experiment is usually associated with a certain redox reaction, and redox reactions between iron oxides, such as wüstite, magnetite, and hematite, have been established as standard oxygen fugacity buffers (Myers and Eugster 1983).

In the past few years, several non-conventional iron oxides, including Fe_4O_5 , Fe_5O_6 , Fe_5O_7 , Fe_7O_9 , $\text{Fe}_{13}\text{O}_{19}$, $\text{Fe}_{25}\text{O}_{32}$, and FeO_2 have been identified experimentally (Bykova et al. 2016; Hu et al. 2016; Lavina et al. 2011; Lavina and Meng 2015; Merlini et al. 2015; Sinmyo et al. 2016). These non-conventional iron oxides form homologous series from several fundamental iron-oxygen polyhedral blocks (Bykova et al. 2016; Guignard and Crichton 2014). Some of these non-conventional iron oxides are quenchable at ambient condition with oxygen fugacities appropriate to the mantle (Guignard and Crichton 2014; Lavina et al. 2011). Besides pure iron oxides, their variants have been identified by substituting Fe^{2+} with Mg^{2+} (Boffa Ballaran et al. 2015; Uenver-Thiele et al. 2017a, 2017b). These non-conventional oxides could be incorporated into the "oxide scale" to constrain the P - T - f_{O_2}

history of petrological assemblages, once their phase diagrams and stability fields are established.

In the study by Uenver-Thiele et al. (2017a), the phase relations of magnesium-iron oxide spinels were explored at high-pressure and high-temperature conditions. Two starting materials, MgFe_2O_4 and $\text{Mg}_{0.5}\text{Fe}_{0.5}^{2+}\text{Fe}_2^{3+}\text{O}_4$, were compressed with multi-anvil presses to 23 GPa and 1500 °C. The oxygen fugacity was carefully controlled in the experiments. The run products were quenched to ambient conditions before further analysis with electron microprobe, backscattered electron image, and X-ray powder diffraction.

Uenver-Thiele et al. first investigated the MgFe_2O_4 system. They found that $\text{MgO}+\text{Fe}_2\text{O}_3$ were the stable assemblage at 1200 °C. At 20 GPa and 1400–1500 °C, the coexistence of $\text{Mg}_2\text{Fe}_2\text{O}_5+\text{Fe}_2\text{O}_3$ and $\text{hp-MgFe}_2\text{O}_4$ phases were observed, and a new phase with the stoichiometry of $\text{Mg}_3\text{Fe}_4\text{O}_9$ were found at 20 GPa, 1300 °C and 23 GPa, 1500 °C, coexisting with Fe_2O_3 and $\text{hp-MgFe}_2\text{O}_4$. Uenver-Thiele et al. suggested that MgO , Fe_2O_3 , $\text{Mg}_2\text{Fe}_2\text{O}_5$, and $\text{Mg}_3\text{Fe}_4\text{O}_9$ could coexist at ~19 GPa and 1200–1250 °C.

As for the $\text{Mg}_{0.5}\text{Fe}_{0.5}^{2+}\text{Fe}_2^{3+}\text{O}_4$ system, periclase was no longer observed in the investigated P - T range. The $\text{Mg}_{0.5}\text{Fe}_{0.5}^{2+}\text{Fe}_2^{3+}\text{O}_4$ spinel broke down to $\text{MgFe}^{2+}\text{Fe}_2^{3+}\text{O}_5 + \text{Fe}_2\text{O}_3$ at 11 GPa and 1000–1600 °C, and the phase boundary was indistinguishable from that for the Fe_3O_4 end-member (Woodland et al. 2012). When the pressure went higher than 16 GPa, a single phase of $\text{hp-Mg}_{0.5}\text{Fe}_{0.5}^{2+}\text{Fe}_2^{3+}\text{O}_4$ became stable. At 15 GPa and 1600 °C, Uenver-Thiele et al. identified a new phase of $\text{Mg}_{1.37}\text{Fe}_{1.63}^{2+}\text{Fe}_4^{3+}\text{O}_9$, another solid solution in the $(\text{Mg}, \text{Fe}^{2+})_3\text{Fe}_4^{3+}\text{O}_9$ series. X-ray powder diffraction suggested that the $(\text{Mg}, \text{Fe}^{2+})_3\text{Fe}_4^{3+}\text{O}_9$ had the same $C2/m$ structure as Fe_3O_9 (Sinmyo et al. 2016).

One interesting application of this study is to constrain the precipitation condition of magnetite inclusions in diamonds. Uenver-Thiele et al. propose an empirical way to determine the precipitation pressure of the quenched magnetite inclusions in diamonds: since the phase diagrams presented in this study suggest that the $\text{hp-(Mg}, \text{Fe}^{2+})\text{Fe}_3^{3+}\text{O}_4$ phase cannot directly transform into the spinel structure, if the precipitated magnetite inclusion has an euhedral morphology, it is likely to come from the partial oxidation of ferropericlase; otherwise, if the magnetite inclusion demonstrates a twinned texture, some kind of precursor phases such as the $(\text{Mg}, \text{Fe}^{2+})_2\text{Fe}_2^{3+}\text{O}_5$ and/or $(\text{Mg}, \text{Fe}^{2+})_3\text{Fe}_4^{3+}\text{O}_9$ phases might be involved in the precipitation process. To sum up, this study has extended our understanding about the high-pressure–high-temperature phase diagrams of magnesium-iron oxides, and provides more detailed constraints on the petrological history of natural diamonds that contain magnesium-iron oxide inclusions.

* E-mail: dzhang@hawaii.edu

REFERENCES CITED

- Boffa Ballaran, T., Uenver-Thiele, L., and Woodland, A.B. (2015) Complete substitution of Fe^{2+} by Mg in Fe_3O_5 : The crystal structure of the $\text{Mg}_2\text{Fe}_2\text{O}_5$ end-member. *American Mineralogist*, 100, 628–632.
- Bykova, E., Dubrovinsky, L., Dubrovinskaia, N., Bykov, M., McCammon, C., Ovsyannikov, S.V., Liermann, H.P., Kupenko, I., Chumakov, A.I., Ruffer, R., Hanfland, M., and Prakapenka, V. (2016) Structural complexity of simple Fe_2O_3 at high pressures and temperatures. *Nature Communications*, 7, 10,661.
- Guignard, J., and Crichton, W.A. (2014) Synthesis and recovery of bulk Fe_3O_5 from magnetite, Fe_3O_4 . A member of a self-similar series of structures for the lower mantle and transition zone. *Mineralogical Magazine*, 78(2), 361–371.
- Hu, Q.Y., Kim, D.Y., Yang, W.G., Yang, L.X., Meng, Y., Zhang, L., and Mao, H.K. (2016) FeO_2 and FeOOH under deep lower-mantle conditions and Earth's oxygen-hydrogen cycles. *Nature*, 534, 241–244.
- Lavina, B., and Meng, Y. (2015) Unraveling the complexity of iron oxides at high pressure and temperature: Synthesis of Fe_5O_6 . *Science Advances*, 1(5), e1400260.
- Lavina, B., Dera, P., Kim, E., Meng, Y., Downs, R.T., Weck, P.F., Sutton, S.R., and Zhao, Y.S. (2011) Discovery of the recoverable high-pressure iron oxide Fe_4O_5 . *Proceedings of the National Academy of Sciences*, 108(42), 17,281–17,285.
- Merlini, M., Hanfland, M., Salamat, A., Petitgirard, S., and Muller, H. (2015) The crystal structures of $\text{Mg}_2\text{Fe}_2\text{C}_2\text{O}_{13}$, with tetrahedrally coordinated carbon, and $\text{Fe}_{13}\text{O}_{19}$, synthesized at deep mantle conditions. *American Mineralogist*, 100, 2001–2004.
- Myers, J., and Eugster, H.P. (1983) The system Fe-Si-O—Oxygen buffer calibrations to 1,500K. *Contributions to Mineralogy and Petrology*, 82(1), 75–90.
- Sinmyo, R., Bykova, E., Ovsyannikov, S.V., McCammon, C., Kupenko, I., Ismailova, L., and Dubrovinsky, L. (2016) Discovery of Fe_2O_5 : a new iron oxide with a complex monoclinic structure. *Scientific Reports*, 6, 32852.
- Uenver-Thiele, L., Woodland, A.B., Boffa Ballaran, T., Miyajima, N., and Frost, D.J. (2017a) Phase relations of Fe-Mg spinels including new high-pressure post-spinel phases and implications for natural samples. *American Mineralogist*, 102, 2054–2064.
- (2017b) Phase relations of MgFe_2O_4 at conditions of the deep upper mantle and transition zone. *American Mineralogist*, 102, 632–642.
- Wirth, R., Dobrzhinetskaya, L., Harte, B., Schreiber, A., and Green, H.W. (2014) High-Fe (Mg,Fe)O inclusion in diamond apparently from the lowermost mantle. *Earth and Planetary Science Letters*, 404, 365–375.
- Woodland, A.B., Frost, D.J., Trots, D.M., Klimm, K., and Mezouar, M. (2012) In situ observation of the breakdown of magnetite (Fe_3O_4) to Fe_2O_3 and hematite at high pressures and temperatures. *American Mineralogist*, 97, 1808–1811.

MANUSCRIPT RECEIVED JUNE 23, 2017
 MANUSCRIPT ACCEPTED JUNE 29, 2017
 MANUSCRIPT HANDLED BY KEITH PUTIRKA

Interested in minerals, rocks, and the processes that form them? Learn about the

Mineralogical Society of America



www.minsocam.org

Mineralogical Society of America 3635 Concorde Pkwy Ste 500 Chantilly VA 20151-1110 USA
 phone: +1 (703) 652-9950 fax: +1 (703) 652-9951

(Sliced and polished, color-zones tourmaline, Plumbago Mining's Newry or Mt. Mica deposit, Newry, Maine, U.S.A.)

Awards
 Short Courses
 Specialist Groups
 Collector's Corner
 Mineralogy 4 Kids
 Member Discounts
 Ask-a-Mineralogist
 MSA-Talk list-serve
 Elements – magazine
 Lectureship Program
 Student Research Grants
 Monographs and Textbooks
 Symposia and Special Sessions
American Mineralogist – journal
Reviews in Mineralogy and Geochemistry

